

Welcome


***“Dark Interactions:
Perspectives from Theory and Experiment”***

***William J. Marciano
Brookhaven National Lab (BNL)
(October 4, 2016)***



BNL is a Multi-Purpose National Lab

- Physics (Nuclear, BES, HEP) Chemistry, Biology, Energy... etc
- Nuclear (RHIC), BES (NSLS II) Major Facilities
- **HEP (ATLAS, Neutrinos [Daya Bay, DUNE...], $g_\mu-2$, Theory...)**



BNL Campus: (RHIC & NSLS Rings)

- 5,320 Acres, ~300 Buildings, ≤ 3000 Employees
- Relativistic Heavy Ion Collider – Last USA Collider!



NSLS II (Newest Light Source) ~\$1 Billion




Photo: Brookhaven National Laboratory


BNL HEP “Physics of the Universe”

Exploring the Frontiers of Energy, Intensity and the Cosmos

- Featured Projects:ATLAS, Daya Bay, DUNE, LSST, BOSS



Bullet Galaxy Cluster Dark Matter Poster



The Dark Boson – A Portal to Dark Matter

What if some dark sector particles interact with one another via a new massive but relatively “light” γ_d (Dark Boson)?

$U(1)_d$ local gauge symmetry of the Dark Sector

Introduced for 1) Sommerfeld Enhancement

- 2) $\gamma_d \rightarrow e^+ e^-$ (source of positrons, **γ -rays**)
- 3) Cosmic Stability of dark matter $U(1)_d$
eg Wimp Number (S. Weinberg)


- 4) Light (< GeV) Dark Matter Abundance

***5) Muon Anomalous Magnetic Moment**
etc.


Can we find direct evidence for such a particle (boson)?

* *Dark Boson g_μ -2 Solution
(Fayet, Pospelov...)*

- **3 Loop Diagram Example**



NA48/2 Updated Bounds on Dark Photon
Simple g_μ -2 discrepancy solution ruled out
Assumes $\text{BR}(Z_d \rightarrow e^+e^-) \sim 1$



BNL to FNAL: The Last Mile

